

FUNDAMENTAL GROUP OF SIMPLE C^* -ALGEBRAS WITH UNIQUE TRACE III

NORIO NAWATA

ABSTRACT. We introduce the fundamental group $\mathcal{F}(A)$ of a simple σ -unital C^* -algebra A with unique (up to scalar multiple) densely defined lower semicontinuous trace. This is a generalization of [29] and [30]. Our definition in this paper makes sense for stably projectionless C^* -algebras. We show that there exist separable stably projectionless C^* -algebras such that their fundamental groups are equal to $\mathbb{R}_+^\times$ by using the classification theorem of Razak [37] and Tsang [40]. This is a contrast to the unital case in [29] and [30]. This study is motivated by the work of Kishimoto and Kumjian in [20].

1. INTRODUCTION

Let M be a factor of type II_1 with a normalized trace τ . Murray and von Neumann introduced the fundamental group $\mathcal{F}(M)$ of M in [28]. They showed that if M is hyperfinite, then $\mathcal{F}(M) = \mathbb{R}_+^\times$. Since then there has been many works on the computation of the fundamental groups. Voiculescu [41] showed that $\mathcal{F}(L(\mathbb{F}_\infty))$ of the group factor of the free group $\mathbb{F}_\infty$ contains the positive rationals and Radulescu proved that $\mathcal{F}(L(\mathbb{F}_\infty)) = \mathbb{R}_+^\times$ in [36]. Connes [7] showed that if G is an ICC group with property (T), then $\mathcal{F}(L(G))$ is a countable group. Popa showed that any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of some factor of type II_1 in [34]. Furthermore Popa and Vaes [35] exhibited a large family $\mathcal{S}$ of subgroups of $\mathbb{R}_+^\times$, containing $\mathbb{R}_+^\times$ itself, all of its countable subgroups, as well as uncountable subgroups with any Hausdorff dimension in $(0, 1)$, such that for each $G \in \mathcal{S}$ there exist many free ergodic measure preserving actions of $\mathbb{F}_\infty$ for which the associated II_1 factor M has the fundamental group equal to G . In our previous paper [29], we introduced the fundamental group $\mathcal{F}(A)$ of a simple unital C^* -algebra A with a normalized trace τ based on the computation of Picard groups by Kodaka [22], [23] and [24]. The fundamental group $\mathcal{F}(A)$ is defined as the set of the numbers $\tau \otimes \text{Tr}(p)$ for some projection $p \in M_n(A)$ such that $pM_n(A)p$ is isomorphic to A . We computed the fundamental groups of several C^* -algebras and showed that any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group of a separable simple unital C^* -algebra with a unique trace in [29] and [30]. Note that the fundamental groups of separable simple unital C^* -algebras are countable.

2000 *Mathematics Subject Classification.* Primary 46L05, Secondary 46L08; 46L35.

Key words and phrases. Fundamental group; Picard group; Hilbert module; Countable basis; Stably projectionless algebra; Dimension function.

In this paper we introduce the fundamental group of a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace. We do not assume that C^* -algebras are unital. In particular, our definition in this paper makes sense for stably projectionless C^* -algebras. Let A be a σ -unital simple C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . The fundamental group $\mathcal{F}(A)$ of A is defined as the set of the numbers $d_\tau(h_1)/d_\tau(h_2)$ for some nonzero positive elements $h_1, h_2 \in A \otimes \mathbb{K}$ such that $\overline{h_1(A \otimes \mathbb{K})h_1}$ is isomorphic to $\overline{h_2(A \otimes \mathbb{K})h_2}$ and $d_\tau(h_2) < \infty$ where d_τ is the dimension function defined by τ . Then the fundamental group $\mathcal{F}(A)$ of A is a multiplicative subgroup of $\mathbb{R}_+^\times$. We show that if A is unital, then our definition in this paper coincides with previous definition in [29] and [30]. Hence if $A \otimes \mathbb{K}$ is separable and has a nonzero projection, then $\mathcal{F}(A)$ is a countable multiplicative subgroup of $\mathbb{R}_+^\times$. By contrast, we show that there exist separable simple stably projectionless C^* -algebras such that their fundamental groups are equal to $\mathbb{R}_+^\times$ by using the classification theorem of Razak [37] and Tsang [40]. This study is motivated by the work of Kishimoto and Kumjian in [20]. (See Example 4.21.)

2. HILBERT C^* -MODULES AND INDUCED TRACES

We say a C^* -algebra A is σ -unital if A has a countable approximate unit. In particular, if A is σ -unital, then there exists a positive element $h \in A$ such that $\{h^{\frac{1}{n}}\}_{n \in \mathbb{N}}$ is an approximate unit. Such a positive element h is called *strict positive* in A . Let $\mathcal{X}$ be a right Hilbert A -module and let $\mathcal{H}(A)$ denote the set of isomorphic classes $[\mathcal{X}]$ of right Hilbert A -modules. (See [26] and [27] for the basic facts on Hilbert modules.) We denote by $L_A(\mathcal{X})$ the algebra of the adjointable operators on $\mathcal{X}$. For $\xi, \eta \in \mathcal{X}$, a "rank one operator" $\Theta_{\xi, \eta}$ is defined by $\Theta_{\xi, \eta}(\zeta) = \xi \langle \eta, \zeta \rangle_A$ for $\zeta \in \mathcal{X}$. We denote by $K_A(\mathcal{X})$ the closure of the linear span of "rank one operators" $\Theta_{\xi, \eta}$ and by $\mathbb{K}$ the C^* -algebra of compact operators on a separable infinite Hilbert space. Let $\mathcal{X}_A$ be a right Hilbert A -module A with the obvious right A -action and $\langle a, b \rangle_A = a^*b$ for $a, b \in A$. Then $K_A(\mathcal{X}_A)$ is isomorphic to A . Hence if A is unital, then $K_A(\mathcal{X}_A) = L_A(\mathcal{X}_A)$. The multiplier algebra, denote by $M(A)$, of a C^* -algebra A is the largest unital C^* -algebra that contains A as an essential ideal. It is unique up to isomorphism over A and isomorphic to $L_A(\mathcal{X}_A)$. Let H_A denote the standard Hilbert module $\{(x_n)_{n \in \mathbb{N}}; x_n \in A, \sum x_n^* x_n \text{ converges in } A\}$ with an A -valued inner product $\langle (x_n)_{n \in \mathbb{N}}, (y_n)_{n \in \mathbb{N}} \rangle = \sum x_n^* y_n$. Then there exists a natural isomorphism ψ of $A \otimes \mathbb{K}$ to $K_A(H_A)$ and ψ can be uniquely extended to an isomorphism $\tilde{\psi}$ of $M(A \otimes \mathbb{K})$ to $L_A(H_A)$. For simplicity of notation, we use the same letter x for $\tilde{\psi}(x)$ where $x \in M(A \otimes \mathbb{K})$.

A finite subset $\{\xi_i\}_{i=1}^n$ of $\mathcal{X}$ is called a *finite basis* if $\eta = \sum_{i=1}^n \xi_i \langle \xi_i, \eta \rangle_A$ for any $\eta \in \mathcal{X}$. More generally, we call a sequence $\{\xi_i\}_{i \in \mathbb{N}} \subseteq \mathcal{X}$ a *countable basis* of $\mathcal{X}$ if $\eta = \sum_{i=1}^\infty \xi_i \langle \xi_i, \eta \rangle_A$ in norm for any $\eta \in \mathcal{X}$, see [17], [18] and [42]. It is also called a standard normalized tight frame as in [12] and [13]. A countable basis $\{\xi_i\}_{i \in \mathbb{N}}$ always converges unconditionally, that is, for any $\eta \in \mathcal{X}$, the net associating $\sum_{i \in F} \xi_i \langle \xi_i, \eta \rangle_A$ to each finite subset $F \subseteq \mathbb{N}$ is

norm converging to η . It is a consequence of the following estimate: for every $\xi \in \mathcal{X}$, $a, b \in K_A(\mathcal{X})$, with $0 \leq a \leq b \leq 1$, $\|\xi - b\xi\|^2 \leq \|\xi\| \|\xi - a\xi\|$.

Proposition 2.1. Let A be a simple C^* -algebra and $\mathcal{X}$ a right Hilbert A -module. Assume that $K_A(\mathcal{X})$ is σ -unital. Then $\mathcal{X}$ has a countable basis.

Proof. Consider a right ideal $\{\Theta_{\xi_0, \zeta} : \zeta \in \mathcal{X}\}$ in $K_A(\mathcal{X})$ for some $\xi_0 \in \mathcal{X}$. Then a similar argument in [4] (Lemma 2.3) proves the proposition. $\square$

Remark 2.2. In general, we need not assume that A is simple. If $K_A(\mathcal{X})$ is σ -unital, then $\mathcal{X}$ has a countable basis. This is an immediate consequence of Kasparov's stabilization trick [19].

Let B be a C^* -algebra. An A - B -equivalence bimodule is an A - B -bimodule $\mathcal{F}$ which is simultaneously a full left Hilbert A -module under a left A -valued inner product ${}_A\langle \cdot, \cdot \rangle$ and a full right Hilbert B -module under a right B -valued inner product $\langle \cdot, \cdot \rangle_B$, satisfying ${}_A\langle \xi, \eta \rangle \zeta = \xi \langle \eta, \zeta \rangle_B$ for any $\xi, \eta, \zeta \in \mathcal{F}$. We say that A is *Morita equivalent* to B if there exists an A - B -equivalence bimodule. It is easy to see that $K_B(\mathcal{F})$ is isomorphic to A . A dual module $\mathcal{F}^*$ of an A - B -equivalence bimodule $\mathcal{F}$ is a set $\{\xi^*; \xi \in \mathcal{F}\}$ with the operations such that $\xi^* + \eta^* = (\xi + \eta)^*$, $\lambda \xi^* = (\overline{\lambda} \xi)^*$, $b \xi^* a = (a^* \xi b^*)^*$, ${}_B\langle \xi^*, \eta^* \rangle = \langle \eta, \xi \rangle_B$ and $\langle \xi^*, \eta^* \rangle_A = {}_A\langle \eta, \xi \rangle$. The bimodule $\mathcal{F}^*$ is a B - A -equivalence bimodule. We refer the reader to [38] and [39] for the basic facts on equivalence bimodules and Morita equivalence.

We review basic facts on the Picard groups of C^* -algebras introduced by Brown, Green and Rieffel in [5]. For A - A -equivalence bimodules $\mathcal{E}_1$ and $\mathcal{E}_2$, we say that $\mathcal{E}_1$ is isomorphic to $\mathcal{E}_2$ as an equivalence bimodule if there exists a $\mathbb{C}$ -linear one-to-one map Φ of $\mathcal{E}_1$ onto $\mathcal{E}_2$ with the properties such that $\Phi(a\xi b) = a\Phi(\xi)b$, ${}_A\langle \Phi(\xi), \Phi(\eta) \rangle = {}_A\langle \xi, \eta \rangle$ and $\langle \Phi(\xi), \Phi(\eta) \rangle_A = \langle \xi, \eta \rangle_A$ for $a, b \in A$, $\xi, \eta \in \mathcal{E}_1$. The set of isomorphic classes $[\mathcal{E}]$ of the A - A -equivalence bimodules $\mathcal{E}$ forms a group under the product defined by $[\mathcal{E}_1][\mathcal{E}_2] = [\mathcal{E}_1 \otimes_A \mathcal{E}_2]$. We call it the *Picard group* of A and denote it by $\text{Pic}(A)$. The identity of $\text{Pic}(A)$ is given by the A - A -bimodule $\mathcal{E} := A$ with ${}_A\langle a_1, a_2 \rangle = a_1 a_2^*$ and $\langle a_1, a_2 \rangle_A = a_1^* a_2$ for $a_1, a_2 \in A$. The inverse element of $[\mathcal{E}]$ in the Picard group of A is the dual module $[\mathcal{E}^*]$. Let α be an automorphism of A , and let $\mathcal{E}_\alpha^A = A$ with the obvious left A -action and the obvious A -valued inner product. We define the right A -action on $\mathcal{E}_\alpha^A$ by $\xi \cdot a = \xi \alpha(a)$ for any $\xi \in \mathcal{E}_\alpha^A$ and $a \in A$, and the right A -valued inner product by $\langle \xi, \eta \rangle_A = \alpha^{-1}(\xi^* \eta)$ for any $\xi, \eta \in \mathcal{E}_\alpha^A$. Then $\mathcal{E}_\alpha^A$ is an A - A -equivalence bimodule. For $\alpha, \beta \in \text{Aut}(A)$, $\mathcal{E}_\alpha^A$ is isomorphic to $\mathcal{E}_\beta^A$ if and only if there exists a unitary $u \in A$ such that $\alpha = \text{ad } u \circ \beta$. Moreover, $\mathcal{E}_\alpha^A \otimes \mathcal{E}_\beta^A$ is isomorphic to $\mathcal{E}_{\alpha \circ \beta}^A$. Hence we obtain an homomorphism ρ_A of $\text{Out}(A)$ to $\text{Pic}(A)$. An A - B -equivalence bimodule $\mathcal{F}$ induces an isomorphism Ψ of $\text{Pic}(A)$ to $\text{Pic}(B)$ by $\Psi([\mathcal{E}]) = [\mathcal{F}^* \otimes \mathcal{E} \otimes \mathcal{F}]$ for $[\mathcal{E}] \in \text{Pic}(A)$. Therefore if A is Morita equivalent to B , then $\text{Pic}(A)$ is isomorphic to $\text{Pic}(B)$.

If A is unital, then any A - B -equivalence bimodule $\mathcal{F}$ is a finitely generated projective B -module as a right module with a finite basis $\{\xi_i\}_{i=1}^n$. Put $p = (\langle \xi_i, \xi_j \rangle_A)_{ij} \in M_n(B)$. Then p is a projection and $\mathcal{F}$ is isomorphic to pB^n as a right Hilbert B -module with an isomorphism of A to $pM_n(B)p$. In the case A is σ -unital, an A - B -equivalence bimodule $\mathcal{F}$ has a countable basis

$\{\xi_i\}_{i \in \mathbb{N}}$ as a right Hilbert B -module by Proposition 2.1. Define p by $p(b_n)_n = (\sum_{m=1}^{\infty} \langle \xi_n, \xi_m \rangle_B b_m)_n$ for $(b_n)_n \in H_B$. Then p is a projection in $L_B(H_B)$ by [16] (Lemma 2.1) and $\sum_{i=1}^n \Theta_{\xi_i, \xi_i} \leq 1_{L_B(H_B)}$ for any $n \in \mathbb{N}$. Therefore $\mathcal{F}$ is isomorphic to pH_B as a right Hilbert module with an isomorphism of A to $p(B \otimes \mathbb{K})p$.

Proposition 2.3. Let A and B be σ -unital simple C^* -algebras and $\mathcal{F}$ an A - B -equivalence bimodule. Then there exists a positive element $h \in A \otimes \mathbb{K}$ such that $\mathcal{F}$ is isomorphic to $\overline{hH_B}$ as a right Hilbert B -module with an isomorphism of A to $\overline{h(B \otimes \mathbb{K})h}$.

Proof. By the discussion above, there exists a projection $p \in M(B \otimes \mathbb{K})$ such that an A - B -equivalence bimodule $\mathcal{F}$ is isomorphic to pH_B as a right Hilbert B -module with an isomorphism of A to $p(B \otimes \mathbb{K})p$. Since $p(B \otimes \mathbb{K})p$ is σ -unital, there exists a strict positive element h in $p(B \otimes \mathbb{K})p$. It is easy to see that $pH_B = \overline{hH_B}$. Therefore $\mathcal{F}$ is isomorphic to $\overline{hH_B}$ as a right Hilbert B -module with an isomorphism of A to $\overline{h(B \otimes \mathbb{K})h}$ as a C^* -algebra. $\square$

Recall that a *trace* on A is a linear map τ on the positive elements of A , with values in $[0, \infty]$ that vanishes at 0 and satisfies the trace identity $\tau(a^*a) = \tau(aa^*)$. If A is simple, then $\tau(a^*a) = 0$ implies $a = 0$. Define $\mathcal{M}_\tau^+ = \{x \geq 0 : \hat{\tau}(x) < \infty\}$ and $\mathcal{M}_\tau = \text{span} \mathcal{M}_\tau^+$. Then $\mathcal{M}_\tau$ is an ideal in A . Every trace τ on A extends a positive linear map on $\mathcal{M}_\tau$. A *normalized trace* is a state on A which is a trace. We say τ is *densely defined* if $\mathcal{M}_\tau$ is a dense ideal in A . In particular, each densely defined trace on A extends a positive linear map on the Pedersen ideal $\text{Ped}(A)$, which is the minimal dense ideal in A . (See [32].) Note that if A is unital, then every densely defined trace is bounded. We review some results about inducing traces from a simple σ -unital C^* -algebra A through a right Hilbert A -module $\mathcal{X}$. See, for example, [6], [9], [14], [25], [31] and [33] for induced traces in several settings. We state the relevant properties in a way that is convenient our purposes, and we include a self-contained proof.

Proposition 2.4. Let A and $\mathcal{X}$ be as above and let τ be a densely defined lower semicontinuous trace. For $x \in K_A(\mathcal{X})_+$ (resp. $L_A(\mathcal{X})_+$), define

$$\text{Tr}_\tau^{\mathcal{X}}(x) := \sum_{i=1}^{\infty} \tau(\langle \xi_i, x\xi_i \rangle_A)$$

where $\{\xi_i\}_{i=1}^{\infty}$ is a countable basis of $\mathcal{X}$. Then $\text{Tr}_\tau^{\mathcal{X}}$ does not depend on the choice of basis and is a densely defined (resp. strictly densely defined) lower semicontinuous trace on $K_A(\mathcal{X})$ (resp. $L_A(\mathcal{X})$).

Proof. Let $\{\xi_i\}_{i \in \mathbb{N}}$ and $\{\zeta_k\}_{k \in \mathbb{N}}$ be countable bases of $\mathcal{X}$. For any positive element $x \in K_A(\mathcal{X})_+$,

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{i=1}^n \tau(\langle \xi_i, x\xi_i \rangle_A) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \tau(\langle \sum_{k=1}^{\infty} \zeta_k \langle \zeta_k, x^{\frac{1}{2}} \xi_i \rangle_A, x^{\frac{1}{2}} \xi_i \rangle_A) \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \tau(\sum_{k=1}^{\infty} \langle x^{\frac{1}{2}} \xi_i, \zeta_k \rangle_A \langle \zeta_k, x^{\frac{1}{2}} \xi_i \rangle_A). \end{aligned}$$

By the lower semicontinuity of τ and $\sum_{i=1}^n \Theta_{\xi_i, \xi_i} \leq 1_{L_A(\mathcal{X})}$,

$$\begin{aligned}
\lim_{n \rightarrow \infty} \sum_{i=1}^n \tau \left(\sum_{k=1}^{\infty} \langle x^{\frac{1}{2}} \xi_i, \zeta_k \rangle_A \langle \zeta_k, x^{\frac{1}{2}} \xi_i \rangle_A \right) &= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sum_{k=1}^{\infty} \tau(\langle x^{\frac{1}{2}} \xi_i, \zeta_k \rangle_A \langle \zeta_k, x^{\frac{1}{2}} \xi_i \rangle_A) \\
&= \lim_{n \rightarrow \infty} \sum_{i=1}^n \sum_{k=1}^{\infty} \tau(\langle x^{\frac{1}{2}} \zeta_k, \xi_i \rangle_A \langle \xi_i, x^{\frac{1}{2}} \zeta_k \rangle_A) \\
&= \lim_{n \rightarrow \infty} \sum_{k=1}^{\infty} \tau \left(\langle x^{\frac{1}{2}} \zeta_k, \sum_{i=1}^n \xi_i \langle \xi_i, x^{\frac{1}{2}} \zeta_k \rangle_A \rangle_A \right) \\
&\leq \lim_{m \rightarrow \infty} \sum_{k=1}^m \tau(\langle \zeta_k, x \zeta_k \rangle_A).
\end{aligned}$$

Therefore $Tr_{\tau}^{\mathcal{X}}$ does not depend on the choice of basis. A similar argument implies that $Tr_{\tau}^{\mathcal{X}}(x^*x) = Tr_{\tau}^{\mathcal{X}}(xx^*)$ for $x \in L_A(\mathcal{X})$.

We shall show that $Tr_{\tau}^{\mathcal{X}}$ is densely defined on $K_A(\mathcal{X})$. Since $K_A(\mathcal{X})$ is simple, it is enough to show that there exists a nonzero element $x \in K_A(\mathcal{X})$ such that $Tr_{\tau}^{\mathcal{X}}(x) < \infty$. There exists a nonzero positive element $a \in A$ such that $\tau(a) < \infty$ because τ is densely defined on A . For any $\eta \in \mathcal{X}$, we have $\langle \eta a^{\frac{1}{2}}, \eta a^{\frac{1}{2}} \rangle_A \leq \|\langle \eta, \eta \rangle_A\| a$, so $\tau(\langle \eta a^{\frac{1}{2}}, \eta a^{\frac{1}{2}} \rangle_A) < \infty$. By the simplicity of A , there exists an element η_0 in $\mathcal{X}$ such that $\eta_0 a^{\frac{1}{2}} \neq 0$. Define $\zeta := \eta_0 a^{\frac{1}{2}}$. Then we have

$$\begin{aligned}
Tr_{\tau}^{\mathcal{X}}(\Theta_{\zeta, \zeta}) &= \sum_{n=1}^{\infty} \tau(\langle \xi_n, \zeta \rangle_A \langle \zeta, \xi_n \rangle_A) \\
&= \lim_{N \rightarrow \infty} \tau \left(\langle \zeta, \sum_{n=1}^N \xi_n \langle \xi_n, \zeta \rangle_A \rangle_A \right) \\
&= \tau(\langle \zeta, \zeta \rangle_A) < \infty
\end{aligned}$$

by the lower semicontinuity of τ . Therefore $Tr_{\tau}^{\mathcal{X}}$ is densely defined on $K_A(\mathcal{X})$. It is easy to see that $Tr_{\tau}^{\mathcal{X}}$ is lower semicontinuous. $\square$

Remark 2.5. Since a right Hilbert A -module $\mathcal{X}$ is a $K_A(\mathcal{X})$ - A -equivalence bimodule, a similar computation in the proof above implies $Tr_{Tr_{\tau}^{\mathcal{X}}}^{\mathcal{X}^*} = \tau$. Therefore there exists a bijective correspondence between densely defined lower semicontinuous traces on A and those on $K_A(\mathcal{X})$.

To simplify notation, we use the same letter τ for the induced trace $Tr_{\tau}^{\mathcal{X}^A}$ on $M(A)$. We denote by $\hat{\tau}$ the induced trace $Tr_{\tau}^{H^A}$ on $M(A \otimes \mathbb{K})$.

3. MULTIPLICATIVE MAPS OF THE PICARD GROUPS TO $\mathbb{R}_+^{\times}$

Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ_A . Define a map $\hat{T}_{\tau_A}$ of $\mathcal{H}(A)$ to $[0, \infty]$ by

$$\hat{T}_{\tau_A}([\mathcal{X}]) := Tr_{\tau_A}^{\mathcal{X}}(1_{L_A(\mathcal{X})}).$$

We see that $\hat{T}_{\tau_A}([\mathcal{X}]) = \sum_{i=1}^{\infty} \tau_A(\langle \xi_i, \xi_i \rangle_A)$ where $\{\xi_i\}_{i=1}^{\infty}$ is a countable basis of $\mathcal{X}$ and it does not depend on the choice of basis (see Proposition 2.4). It is easily seen that $\hat{T}_{\tau_A}$ is well-defined. We shall compute $\hat{T}_{\tau_A}(\overline{[hH_A]})$ where h is

a positive element in $A \otimes \mathbb{K}$. Let $d_{\tau_A}(h) = \lim_{n \rightarrow \infty} \hat{\tau}_A(h^{\frac{1}{n}})$ for $h \in (A \otimes \mathbb{K})_+$. Then d_{τ_A} is a dimension function. (See, for example, [1], [2] and [8].)

Proposition 3.1. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ_A and h a positive element in $A \otimes \mathbb{K}$. Then $\hat{T}_{\tau_A}([\overline{hH_A}]) = d_{\tau_A}(h)$.

Proof. We may assume that $\|h\| \leq 1$. Then $\{h^{\frac{1}{n}}\}_{n \in \mathbb{N}}$ is an increasing approximate unit for $K_A(\overline{hH_A})$ and $\lim_{n \rightarrow \infty} h^{\frac{1}{n}}\xi = \xi$ for any $\xi \in \overline{hH_A}$. Let $\{\xi_i\}_{i \in \mathbb{N}}$ be a basis of $\overline{hH_A}$ and $\{\eta_j\}_{j \in \mathbb{N}}$ a basis of H_A . By the lower semicontinuity of τ_A and $\langle \xi_i, h^{\frac{1}{n}}\xi_i \rangle_A \leq \langle \xi_i, h^{\frac{1}{n+1}}\xi_i \rangle_A$, we have

$$\begin{aligned} \lim_{n \rightarrow \infty} \sum_{i=1}^{\infty} \tau_A(\langle \xi_i, h^{\frac{1}{n}}\xi_i \rangle_A) &= \sum_{i=1}^{\infty} \lim_{n \rightarrow \infty} \tau_A(\langle \xi_i, h^{\frac{1}{n}}\xi_i \rangle_A) \\ &= \sum_{i=1}^{\infty} \tau_A(\langle \xi_i, \xi_i \rangle_A) = \hat{T}_{\tau_A}([\overline{hH_A}]). \end{aligned}$$

In a similar way, we have

$$\begin{aligned} \sum_{i=1}^{\infty} \tau_A(\langle \xi_i, h^{\frac{1}{n}}\xi_i \rangle_A) &= \sum_{i=1}^{\infty} \tau_A(\langle \xi_i, h^{\frac{1}{2n}} \sum_{j=1}^{\infty} \eta_j \langle \eta_j, h^{\frac{1}{2n}}\xi_i \rangle_A \rangle_A) \\ &= \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tau_A(\langle \xi_i, h^{\frac{1}{2n}}\eta_j \rangle_A \langle h^{\frac{1}{2n}}\eta_j, \xi_i \rangle_A) \\ &= \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tau_A(\langle h^{\frac{1}{2n}}\eta_j, \xi_i \rangle_A \langle \xi_i, h^{\frac{1}{2n}}\eta_j \rangle_A) \\ &= \sum_{j=1}^{\infty} \sum_{i=1}^{\infty} \tau_A(\langle h^{\frac{1}{2n}}\eta_j, \xi_i \rangle_A \langle \xi_i, h^{\frac{1}{2n}}\eta_j \rangle_A) \\ &= \sum_{j=1}^{\infty} \tau_A(\langle \eta_j, h^{\frac{1}{n}}\eta_j \rangle_A) = \hat{\tau}_A(h^{\frac{1}{n}}). \end{aligned}$$

Therefore $\hat{T}_{\tau_A}([\overline{hH_A}]) = d_{\tau_A}(h)$. $\square$

Remark 3.2. Let p be a projection in $M(A \otimes \mathbb{K})$. Then it is easy to see that $\hat{T}_{\tau_A}([pH_A]) = \hat{\tau}_A(p)$.

The following proposition is a generalization of Proposition 2.1 in [30].

Proposition 3.3. Let A and B be simple σ -unital C^* -algebras with unique (up to scalar multiple) densely defined lower semicontinuous traces τ_A and τ_B respectively. Assume that $\tau_A(1_{M(A)}) = 1$, that is, τ_A is a normalized trace. Then for every right Hilbert A -module $\mathcal{X}$ and every A - B -equivalence bimodule $\mathcal{F}$,

$$\hat{T}_{\tau_B}([\mathcal{X} \otimes \mathcal{F}]) = \hat{T}_{\tau_A}([\mathcal{X}])\hat{T}_{\tau_B}([\mathcal{F}]).$$

Proof. Let $\{\xi_i\}_{i \in \mathbb{N}}$ be a countable basis of $\mathcal{X}$ and $\{\eta_j\}_{j \in \mathbb{N}}$ a countable basis of $\mathcal{F}$ as a right Hilbert B -module. Then $\{\xi_i \otimes \eta_j\}_{i,j \in \mathbb{N}}$ is a countable basis

of $\mathcal{X} \otimes \mathcal{F}$ as a right Hilbert A -module. By $\tau_B(\langle \xi_i \otimes \eta_j, \xi_i \otimes \eta_j \rangle_B) \geq 0$, we have

$$\begin{aligned} \hat{T}_{\tau_B}([\mathcal{X} \otimes \mathcal{F}]) &= \sum_{i,j=1}^{\infty} \tau_B(\langle \xi_i \otimes \eta_j, \xi_i \otimes \eta_j \rangle_B) = \sum_{i=1}^{\infty} \sum_{j=1}^{\infty} \tau_B(\langle \eta_j, \langle \xi_i, \xi_i \rangle_A \eta_j \rangle_B) \\ &= \sum_{i=1}^{\infty} \text{Tr}_{\tau_B}^{\mathcal{F}}(\langle \xi_i, \xi_i \rangle_A). \end{aligned}$$

Since $\text{Tr}_{\tau_B}^{\mathcal{F}}$ is a densely defined lower semicontinuous trace on A , there exists $\lambda \in \mathbb{R}_+^{\times}$ such that $\text{Tr}_{\tau_B}^{\mathcal{F}} = \lambda \tau_A$. The assumption $\tau_A(1_{M(A)}) = 1$ implies $\lambda = \text{Tr}_{\tau_B}^{\mathcal{F}}(1_{M(A)})$. Therefore

$$\sum_{i=1}^{\infty} \text{Tr}_{\tau_B}^{\mathcal{F}}(\langle \xi_i, \xi_i \rangle_A) = \sum_{i=1}^{\infty} \text{Tr}_{\tau_B}^{\mathcal{F}}(1_{M(A)}) \tau_A(\langle \xi_i, \xi_i \rangle_A) = \hat{T}_{\tau_A}([\mathcal{X}]) \hat{T}_{\tau_B}([\mathcal{F}]).$$

□

We shall consider the multiplicative map of the Picard group to $\mathbb{R}_+^{\times}$.

Proposition 3.4. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ_A . Assume that $\mathcal{X}$ is a nonzero right Hilbert A -module such that $\hat{T}_{\tau_A}([\mathcal{X}]) < \infty$. Define a map $T_{\mathcal{X}}$ of $\text{Pic}(K_A(\mathcal{X}))$ to $\mathbb{R}_+^{\times}$ by

$$T_{\mathcal{X}}([\mathcal{E}]) := \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}])} \hat{T}_{\tau_A}([\mathcal{E} \otimes \mathcal{X}])$$

for $[\mathcal{E}] \in \text{Pic}(K_A(\mathcal{X}))$. Then $T_{\mathcal{X}}$ is well-defined and independent on the choice of trace. Moreover $T_{\mathcal{X}}$ is a multiplicative map.

Proof. If $K_A(\mathcal{X})$ - $K_A(\mathcal{X})$ equivalence bimodule $\mathcal{E}'$ is isomorphic to $\mathcal{E}$, then $\mathcal{E}' \otimes \mathcal{X}$ is isomorphic to $\mathcal{E} \otimes \mathcal{X}$. Hence $T_{\mathcal{X}}([\mathcal{E}']) = T_{\mathcal{X}}([\mathcal{E}])$. A similar computation in the proof of Proposition 3.3 shows

$$T_{\mathcal{X}}([\mathcal{E}]) = \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}])} \hat{T}_{\tau_A}([\mathcal{E}]) = \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}])} \text{Tr}_{\tau_A}^{\mathcal{E}}(1_{L_A(\mathcal{E})}).$$

Since $\mathcal{E}$ is a $K_A(\mathcal{X})$ - $K_A(\mathcal{X})$ -equivalence bimodule, $K_A(\mathcal{E})$ is isomorphic to $K_A(\mathcal{X})$. The uniqueness of the trace on $K_A(\mathcal{X})$ implies

$$\text{Tr}_{\tau_A}^{\mathcal{E}}(1_{L_A(\mathcal{E})}) = \lambda \text{Tr}_{\tau_A}^{\mathcal{X}}(1_{L_A(\mathcal{X})}) = \lambda \hat{T}_{\tau_A}([\mathcal{X}]) < \infty$$

for some $\lambda \in \mathbb{R}_+^{\times}$. Therefore $T_{\mathcal{X}}$ is well-defined. Define $\tau' := \frac{\text{Tr}_{\tau_A}^{\mathcal{X}}}{\hat{T}_{\tau_A}([\mathcal{X}])}$. Then τ' is a normalized trace on $K_A(\mathcal{X})$. By proposition 3.3,

$$\begin{aligned} T_{\mathcal{X}}([\mathcal{E}][\mathcal{E}']) &= \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}])} \hat{T}_{\tau_A}([\mathcal{E} \otimes \mathcal{E}' \otimes \mathcal{X}]) = \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}])} \hat{T}_{\tau_A}([\mathcal{E} \otimes \mathcal{E}']) \\ &= \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}])} \hat{T}_{\tau'}([\mathcal{E}]) \hat{T}_{\tau_A}([\mathcal{E}']) \\ &= T_{\mathcal{X}}([\mathcal{E}]) T_{\mathcal{X}}([\mathcal{E}']). \end{aligned}$$

□

4. FUNDAMENTAL GROUPS

Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ , and let h_0 be a nonzero positive element in $A \otimes \mathbb{K}$ with $d_\tau(h_0) < \infty$. Put

$$\mathcal{F}_{h_0}(A) := \left\{ d_\tau(h)/d_\tau(h_0) \in \mathbb{R}_+^\times \mid \begin{array}{l} h \text{ is a positive element in } A \otimes \mathbb{K} \text{ such that} \\ \overline{h(A \otimes \mathbb{K})h} \cong \overline{h_0(A \otimes \mathbb{K})h_0} \end{array} \right\}$$

Lemma 4.1. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ and h_0 a nonzero positive element in $A \otimes \mathbb{K}$ such that $d_\tau(h_0) < \infty$. Then $\mathcal{F}_{h_0}(A)$ is a multiplicative subgroup of $\mathbb{R}_+^\times$.

Proof. Put $\mathcal{X} = \overline{h_0 H_A}$. It is enough to show that $\mathcal{F}_{h_0}(A) = \text{Im}(T_{\mathcal{X}})$. Let $\mathcal{E}$ be a $K_A(\mathcal{X})$ - $K_A(\mathcal{X})$ -equivalence bimodule. Then there exists a positive element $h \in A \otimes \mathbb{K}$ such that $\mathcal{E} \otimes \mathcal{X}$ is isomorphic to $\overline{h H_A}$ as a right Hilbert A -module with an isomorphism of $K_A(\mathcal{X})$ to $\overline{h(A \otimes \mathbb{K})h}$ by Proposition 2.3. Since $K_A(\mathcal{X})$ is isomorphic to $\overline{h_0(A \otimes \mathbb{K})h_0}$ and we have $T_{\mathcal{X}}([\mathcal{E}]) = d_\tau(h)/d_\tau(h_0)$ by Proposition 3.1, $\text{Im}(T_{\mathcal{X}}) \subset \mathcal{F}_{h_0}(A)$. Conversely let h be a positive element in $A \otimes \mathbb{K}$ such that $\overline{h(A \otimes \mathbb{K})h}$ is isomorphic to $\overline{h_0(A \otimes \mathbb{K})h_0}$. Since A is simple and $\overline{h(A \otimes \mathbb{K})h}$ is isomorphic to $K_A(\mathcal{X})$, $\mathcal{E} := \overline{h H_A} \otimes \mathcal{X}^*$ is a $K_A(\mathcal{X})$ - $K_A(\mathcal{X})$ -equivalence bimodule. By Proposition 3.1, $T_{\mathcal{X}}([\mathcal{E}]) = \frac{1}{\hat{T}_{\tau_A}([\mathcal{X}]}) \hat{T}_{\tau_A}([\overline{h H_A}]) = d_\tau(h)/d_\tau(h_0)$. Therefore $\mathcal{F}_{h_0}(A) \subset \text{Im}(T_{\mathcal{X}})$. $\square$

Lemma 4.2. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . Assume that h_0 and h_1 are nonzero positive elements in $A \otimes \mathbb{K}$ such that $d_\tau(h_0), d_\tau(h_1) < \infty$. Then $\mathcal{F}_{h_0}(A) = \mathcal{F}_{h_1}(A)$.

Proof. Let $\mathcal{F} := \overline{h_0 H_A} \otimes (\overline{h_1 H_A})^*$. Then $\mathcal{F}$ is a $\overline{h_0(A \otimes \mathbb{K})h_0}$ - $\overline{h_1(A \otimes \mathbb{K})h_1}$ -equivalence bimodule by the simplicity of A , and $\mathcal{F}$ induces an isomorphism Ψ of $\text{Pic}(\overline{h_0(A \otimes \mathbb{K})h_0})$ to $\text{Pic}(\overline{h_1(A \otimes \mathbb{K})h_1})$ such that $\Psi([\mathcal{E}]) = [\mathcal{F}^* \otimes \mathcal{E} \otimes \mathcal{F}]$ for $[\mathcal{E}] \in \text{Pic}(\overline{h_0(A \otimes \mathbb{K})h_0})$. By Proposition 3.3, $T_{\overline{h_1 H_A}}(\Psi([\mathcal{E}])) = T_{\overline{h_0 H_A}}([\mathcal{E}])$. Therefore $\mathcal{F}_{h_0}(A) = \mathcal{F}_{h_1}(A)$ by the proof of Lemma 4.1. $\square$

Put

$$\mathcal{F}(A) := \left\{ d_\tau(h_1)/d_\tau(h_2) \in \mathbb{R}_+^\times \mid \begin{array}{l} h_1 \text{ and } h_2 \text{ are nonzero positive elements in} \\ A \otimes \mathbb{K} \text{ such that} \\ \overline{h_1(A \otimes \mathbb{K})h_1} \cong \overline{h_2(A \otimes \mathbb{K})h_2}, d_\tau(h_2) < \infty \end{array} \right\}.$$

Theorem 4.3. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . Then $\mathcal{F}(A)$ is a multiplicative subgroup of $\mathbb{R}_+^\times$.

Proof. Let h_0 be a nonzero positive element in $\text{Ped}(A \otimes \mathbb{K})$. By [32] (Proposition 5.6.2), $\overline{h_0(A \otimes \mathbb{K})h_0}$ is contained in $\text{Ped}(A \otimes \mathbb{K})$. Since $\hat{\tau}$ is densely defined, $\text{Ped}(A \otimes \mathbb{K}) \subset \mathcal{M}_{\hat{\tau}}$. Therefore $\hat{\tau}$ is bounded on $\overline{h_0(A \otimes \mathbb{K})h_0}$ and hence $d_\tau(h_0) < \infty$. Lemma 4.2 implies $\cup_{d_\tau(h) < \infty} \mathcal{F}_h(A) = \mathcal{F}_{h_0}(A)$. It is clear that $\mathcal{F}(A) = \cup_{d_\tau(h) < \infty} \mathcal{F}_h(A)$. Consequently $\mathcal{F}(A)$ is a multiplicative subgroup of $\mathbb{R}_+^\times$ by Lemma 4.1. $\square$

Definition 4.4. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . We call $\mathcal{F}(A)$ the fundamental group of A , which is a multiplicative subgroup of $\mathbb{R}_+^\times$.

Remark 4.5. It is easy to see that $\mathcal{F}(A)$ is equal to the set

$$\left\{ \hat{\tau}(p)/\hat{\tau}(q) \in \mathbb{R}_+^\times \mid \begin{array}{l} p \text{ and } q \text{ are nonzero projections in} \\ M(A \otimes \mathbb{K}) \text{ such that} \\ p(A \otimes \mathbb{K})p \cong q(A \otimes \mathbb{K})q, \hat{\tau}(q) < \infty \end{array} \right\}.$$

Remark 4.6. If a unique densely defined lower semicontinuous trace τ is a normalized trace, then $\mathcal{F}(A)$ is equal to the set

$$\{d_\tau(h) \in \mathbb{R}_+^\times \mid h \text{ is a positive element in } A \otimes \mathbb{K} \text{ such that } A \cong \overline{h(A \otimes \mathbb{K})h}\}.$$

Note that there exists a σ -unital simple C^* -algebra with a unique normalized trace τ , which has a densely defined lower semicontinuous trace that is not a scalar multiple of τ . For example, let A be an AF-algebra such that $K_0(A) = \mathbb{Z}[\frac{1}{2}] \oplus \mathbb{Z}[\frac{1}{2}]$, $K_0(A)_+ = \{(q, r) \in K_0(A) : q > 0, r > 0\} \cup \{(0, 0)\}$ and $\Sigma(A) = \{(q, r) \in K_0(A)_+ : q > 0, 0 < r < 1\} \cup \{(0, 0)\}$. Then A is such a C^* -algebra.

The following corollary is shown by a similar argument of Lemma 4.2.

Corollary 4.7. Let A and B be simple σ -unital C^* -algebras with unique (up to scalar multiple) densely defined lower semicontinuous traces. If A is Morita equivalent to B , then $\mathcal{F}(A) = \mathcal{F}(B)$.

We shall show that if A is unital, then Definition 4.4 coincides with previous definition in [29] and [30].

Proposition 4.8. Let A be a unital simple C^* -algebra with a unique normalized trace τ . Then

$$\mathcal{F}(A) = \{\tau \otimes Tr(p) \in \mathbb{R}_+^\times \mid p \text{ is a projection in } M_n(A) \text{ such that } pM_n(A)p \cong A\}$$

where Tr is the usual unnormalized trace on $M_n(\mathbb{C})$.

Proof. Let $\mathcal{X}_A$ be a right Hilbert A -module A with the obvious right A -action and $\langle a, b \rangle_A = a^*b$ for $a, b \in A$. Since τ is a normalized trace, $\hat{T}_\tau([\mathcal{X}_A]) = 1$. By the proof of Lemma 4.1 and Lemma 4.2, $\mathcal{F}(A) = \mathcal{F}_{1 \otimes e_{11}}(A) = \hat{T}_\tau(\text{Pic}(A))$ where e_{11} is a rank one projection in $\mathbb{K}$. A similar argument in [29] (Theorem 3.1) shows $\hat{T}_\tau(\text{Pic}(A)) = \{\tau \otimes Tr(p) \in \mathbb{R}_+^\times \mid p \text{ is a projection in } M_n(A) \text{ such that } pM_n(A)p \cong A\}$ because every A - A -equivalence bimodule has a finite basis. $\square$

We showed that K-theoretical obstruction enables us to compute fundamental groups easily in the case A is unital [29]. Therefore if $A \otimes \mathbb{K}$ has a nonzero projection, we can compute fundamental groups easily by K-theoretical obstruction. We denote by τ_* the map $K_0(A) \rightarrow \mathbb{R}$ induced by a trace τ on A .

Definition 4.9. Let E be an additive subgroup of $\mathbb{R}$ containing $\mathbb{Z}$. Then an *inner multiplier group* $IM(E)$ of E is defined by

$$IM(E) = \{t \in \mathbb{R}^\times \mid t \in E, t^{-1} \in E, \text{ and } tE = E\}.$$

Then $IM(E)$ is a multiplicative subgroup of $\mathbb{R}^\times$. We call $IM_+(E) := IM(E) \cap \mathbb{R}_+$ the *positive inner multiplier group* of E , which is a multiplicative subgroup of $\mathbb{R}_+^\times$.

Corollary 4.10. Let A be a separable simple C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . Assume that $A \otimes \mathbb{K}$ has a nonzero projection. Then $\mathcal{F}(A)$ is countable. Moreover $\tau_*(K_0(A))$ is a $\mathbb{Z}[\mathcal{F}(A)]$ -module and $\mathcal{F}(A) \subset IM_+(\tau_*(K_0(A)))$.

Proof. Let p be a nonzero projection in $A \otimes \mathbb{K}$. Corollary 4.7 implies $\mathcal{F}(A) = \mathcal{F}(p(A \otimes \mathbb{K})p)$. Since $p(A \otimes \mathbb{K})p$ is a separable unital C^* -algebra, [29] (Proposition 3.7) and Proposition 4.8 prove the corollary. $\square$

Example 4.11. Let $\mathbb{F}_n$ be a non-abelian free group with $n \geq 2$ generators. Then $C_r^*(\mathbb{F}_n)$ is a unital simple C^* -algebra with a unique normalized trace. Since $K_0(C_r^*(\mathbb{F}_n)) \cong \mathbb{Z}$, $\mathcal{F}(C_r^*(\mathbb{F}_n)) = \{1\}$. This implies that for positive elements $h_1, h_2 \in C_r^*(\mathbb{F}_n)$ if $\overline{h_1 C_r^*(\mathbb{F}_n) h_1}$ is isomorphic to $\overline{h_2 C_r^*(\mathbb{F}_n) h_2}$, then $d_\tau(h_1) = d_\tau(h_2)$.

Example 4.12. Let p be a prime number. Consider a tensor product algebra of a UHF algebra and the compact operators $A = M_{p^\infty} \otimes \mathbb{K}$. Then $\mathcal{F}(A) = \{p^n : n \in \mathbb{Z}\}$.

Remark 4.13. Any countable subgroup of $\mathbb{R}_+^\times$ can be realized as the fundamental group $\mathcal{F}(A)$ of a separable simple unital C^* -algebra A with a unique trace. (See [30].)

We show that there exist separable simple stably projectionless C^* -algebras such that their fundamental groups are equal to $\mathbb{R}_+^\times$. This is a contrast to the unital case. Recall the building blocks that are considered by Razak [37] and Tsang [40]. These algebras are subhomogeneous algebras obtained by generalized mapping torus construction as in [10] and [11]. For a pair of natural numbers (n, m) with n dividing m ($m > n$), let ρ_0 and ρ_1 be homomorphisms from $M_n(\mathbb{C})$ to $M_m(\mathbb{C})$, which having multiplicities $\frac{m}{n} - 1$ and $\frac{m}{n}$ respectively. Define

$$A(n, m) = \{f \in M_m(C([0, 1])) : f(0) = \rho_0(c), f(1) = \rho_1(c), c \in M_n(\mathbb{C})\}.$$

Note that we may assume that the homomorphism ρ_0 maps $M_n(\mathbb{C})$ into diagonal block matrices in $M_m(\mathbb{C})$ with $\frac{m}{n} - 1$ identical blocks and one zero block, on the other hand, the homomorphism ρ_1 yields matrices with $\frac{m}{n}$ identical blocks. The building block $A(n, m)$ has the following properties. (See, for example, [32] and [37].)

Proposition 4.14. We have the following.

- (i) Every primitive ideal of $A(n, m)$ is the kernel of some point evaluation. Therefore the primitive ideal space of $A(n, m)$ is homeomorphic to $\mathbb{T}$.
- (ii) The Pedersen ideal of $A(n, m)$ is $A(n, m)$. Therefore every densely defined lower semicontinuous trace on $A(n, m)$ is bounded.
- (iii) For any bounded trace τ on $A(n, m)$, there exists a measure μ on $\mathbb{T}$ such that $\tau(f) = \int_{\mathbb{T}} (\frac{m-n}{m})^t \text{Tr}(f(t)) d\mu(t)$ for any $f \in A(n, m)$.

Fix an irrational $\theta \in [0, 1] \setminus \mathbb{Q}$. For any $n \in \mathbb{N}$, define an injective homomorphism ϕ_n of $A(3^n, 2 \cdot 3^n)$ to $A(3^{n+1}, 2 \cdot 3^{n+1})$ by

$$(\phi_n(f))(t) = \begin{cases} u_t \begin{pmatrix} f(t) & 0 & 0 \\ 0 & f(t+\theta) & 0 \\ 0 & 0 & 0 \end{pmatrix} u_t^* & 0 \leq t \leq 1-\theta \\ w_t \begin{pmatrix} f(t) & 0 & 0 \\ 0 & f(t+\theta-1) & 0 \\ 0 & 0 & f(t+\theta-1) \end{pmatrix} w_t^* & 1-\theta \leq t \leq 1 \end{cases}$$

where u_t and w_t are suitable continuous paths in $U(M_{2 \cdot 3^{n+1}}(\mathbb{C}))$. We denote by $\phi_{n,m}$ a homomorphism $\phi_{m-1} \circ \cdots \circ \phi_n$ from $A(3^n, 2 \cdot 3^n)$ to $A(3^m, 2 \cdot 3^m)$. Let $\mathcal{O} = \varinjlim (A(3^n, 2 \cdot 3^n), \phi_{n,m})$.

Lemma 4.15. With notation as above $\mathcal{O} = \varinjlim (A(3^n, 2 \cdot 3^n), \phi_{n,m})$ is a separable simple stably projectionless C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous unbounded trace.

Proof. Let J be a proper two-sided closed ideal of $\mathcal{O}$, and let $J_n = \phi_{n,\infty}^{-1}(J \cap \phi_{n,\infty}(A(3^n, 2 \cdot 3^n)))$. Then J_n is a two-sided closed ideal of $A(3^n, 2 \cdot 3^n)$, and denote by F_n the corresponding closed set in $\mathbb{T}$. (See Proposition 4.14.) Since $\phi_{n,m}$ is injective, $J = \varinjlim (J_n, \phi_{n,m})$ and for n sufficiently large, J_n is a proper two-sided closed ideal of $A(3^n, 2 \cdot 3^n)$, that is, F_n is not empty. Put $\gamma([t]) = [t+\theta]$ for any $[t] \in \mathbb{T}$. For any natural number k , we see that $F_n = F_{n+k} \cup \gamma^{-1}(F_{n+k}) \cup \cdots \cup \gamma^{-k}(F_{n+k})$ by the construction of $\phi_{n,m}$ and $J_n = \phi_{n,n+k}^{-1}(J \cap \phi_{n,n+k}(A(3^n, 2 \cdot 3^n)))$. A same argument in [3](the last part of the proof of Proposition 1.3) shows that $\mathcal{O}$ is simple because γ is minimal homeomorphism on $\mathbb{T}$.

Define $\tau_n(f) = \frac{1}{(1+2^\theta)^n} \int_{\mathbb{T}} (\frac{1}{2})^t \text{Tr}(f(t)) d\mu(t)$ where μ is a normalized Haar measure on $\mathbb{T}$ and Tr is the usual unnormalized trace on $M_{2 \cdot 3^n}(\mathbb{C})$. Then $\tau_n = \tau_{n+1} \circ \phi_n$, and hence there exists a densely defined lower semicontinuous trace τ on $\mathcal{O}$. Note that τ is unbounded trace since $\|\tau_n\| = \frac{2 \cdot 3^n}{(1+2^\theta)^n}$.

We shall show that the uniqueness of τ . Let τ' be a densely defined lower semicontinuous trace on $\mathcal{O}$. It is easy to see that $\tau'|_{A(3^n, 2 \cdot 3^n)}$ is densely defined lower semicontinuous trace on $A(3^n, 2 \cdot 3^n)$. Proposition 4.14 implies that for any $n \in \mathbb{N}$ there exists a measure ν_n on $\mathbb{T}$ such that $\tau'|_{A(3^n, 2 \cdot 3^n)}(f) = \frac{1}{(1+2^\theta)^n} \int_{\mathbb{T}} (\frac{1}{2})^t \text{Tr}(f(t)) d\nu_n(t)$. By a compatibility condition, we have

$$\int_{\mathbb{T}} (\frac{1}{2})^t \text{Tr}(f(t)) d\nu_n(t) = \frac{1}{1+2^\theta} \int_{\mathbb{T}} (\frac{1}{2})^t (\text{Tr}(f(t)) + g(t)) d\nu_{n+1}(t)$$

where

$$g(t) = \begin{cases} \text{Tr}(f(t+\theta)) & 0 \leq t \leq 1-\theta \\ 2\text{Tr}(f(t+\theta-1)) & 1-\theta \leq t \leq 1 \end{cases}$$

for any $f \in A(3^n, 2 \cdot 3^n)$. Therefore for any $h \in C(\mathbb{T})$, we have

$$\int_{\mathbb{T}} h(t) d\nu_n(t) = \frac{1}{1+2^\theta} \int_{\mathbb{T}} h(t) + 2^\theta h(t+\theta) d\nu_{n+1}(t).$$

In a same way in [20](the last part of the proof of Theorem 2.4), we see that ν_n is Haar measure on $\mathbb{T}$ by this condition. Consequently there exists a positive number λ such that $\tau' = \lambda\tau$. $\square$

The following lemma is an immediate consequence of the classification theorem of Razak [37] and Tsang [40](Theorem 3.1).

Lemma 4.16. Let A be an simple separable AF algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace. Then $A \otimes \mathcal{O}$ is isomorphic to $\mathcal{O}$.

Theorem 4.17. There exist a separable simple stably projectionless nuclear C^* -algebra and non-nuclear C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace such that their fundamental groups are equal to $\mathbb{R}_+^\times$.

Proof. For any $\lambda \in \mathbb{R}_+^\times$, there exists a separable unital simple AF algebra A_λ with a unique trace such that $\lambda \in \mathcal{F}(A_\lambda)$ by Corollary 3.16 in [29]. Lemma 4.16 implies $\lambda \in \mathcal{F}(\mathcal{O})$. Therefore $\mathcal{F}(\mathcal{O}) = \mathbb{R}_+^\times$. Let $\mathbb{F}_n$ be a non-abelian free group with $n \geq 2$ generators. Then $\mathcal{O} \otimes C_r^*(\mathbb{F}_n)$ is a separable stably projectionless non-nuclear C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace such that $\mathcal{F}(\mathcal{O} \otimes C_r^*(\mathbb{F}_n)) = \mathbb{R}_+^\times$. $\square$

Remark 4.18. Let h be a nonzero positive element in the Pedersen ideal of $\mathcal{O}$. Then $\overline{h\mathcal{O}h}$ is a separable stably projectionless C^* -algebra with a unique normalized trace such that $\mathcal{F}(\overline{h\mathcal{O}h}) = \mathbb{R}_+^\times$.

Remark 4.19. Recently, Jacelon [15] construct a simple, nuclear, stably projectionless C^* -algebra W with a unique normalized trace, which shares some of the important properties of the Cuntz algebra $\mathcal{O}_2$. This C^* -algebra is an inductive limit of building blocks $A(n, m)$. Hence $W \otimes \mathbb{K}$ is isomorphic to $\mathcal{O}$ by the classification theorem of Razak [37]. Therefore Corollary 4.7 and Theorem 4.17 imply $\mathcal{F}(W) = \mathbb{R}_+^\times$.

Recall that the fundamental group of a II_1 -factor M is equal to the set of trace-scaling constants for automorphisms of $M \otimes B(\mathcal{H})$. We have a similar fact as discussed by Kodaka in [24]. We define the set of trace-scaling constants for automorphisms:

$$\mathfrak{S}(A) := \{\lambda \in \mathbb{R}_+^\times \mid \hat{\tau} \circ \alpha = \lambda \hat{\tau} \text{ for some } \alpha \in \text{Aut}(A \otimes K(\mathcal{H}))\}.$$

Proposition 4.20. Let A be a simple σ -unital C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . Then $\mathcal{F}(A) = \mathfrak{S}(A)$.

Proof. There exists a nonzero projection p in $M(A \otimes \mathbb{K})$ such that $\hat{\tau}(p) < \infty$ by a similar argument in the proof of Theorem 4.3. Let $\lambda \in \mathfrak{S}(A)$, then there exists an automorphism of $A \otimes \mathbb{K}$ such that $\hat{\tau} \circ \alpha(x) = \lambda \hat{\tau}(x)$ for $x \in \mathcal{M}_{\hat{\tau}}$. There exists an automorphism $\tilde{\alpha}$ of $M(A \otimes \mathbb{K})$ such that $\tilde{\alpha}(x) = \alpha(x)$ for $x \in A \otimes \mathbb{K}$. It is clear that $p(A \otimes \mathbb{K})p$ is isomorphic to $\tilde{\alpha}(p)(A \otimes \mathbb{K})\tilde{\alpha}(p)$. We have that $\hat{\tau}(\tilde{\alpha}(p))/\hat{\tau}(p) = \lambda$. Therefore $\lambda \in \mathcal{F}(A)$ by Remark 4.5.

Conversely, let $\lambda \in \mathcal{F}(A)$. There exist projections p and q in $M(A \otimes \mathbb{K})$ such that $p(A \otimes \mathbb{K})p$ is isomorphic to $q(A \otimes \mathbb{K})q$ and $\lambda = \hat{\tau}(p)/\hat{\tau}(q)$. We denote by ϕ an isomorphism of $p(A \otimes \mathbb{K})p$ to $q(A \otimes \mathbb{K})q$. Since p and q are full projections, there exist partial isometries w_1 and w_2 in $(A \otimes \mathbb{K}) \otimes \mathbb{K}$ such that $w_1^*w_1 = I \otimes I$, $w_1w_1^* = p \otimes I$, $w_2^*w_2 = I \otimes I$ and $w_2w_2^* = q \otimes I$ by

Brown [4]. Let $\psi : A \otimes \mathbb{K} \otimes \mathbb{K} \rightarrow A \otimes \mathbb{K}$ be an isomorphism which induces the identity on the K_0 -group. Define $\alpha = \psi \circ (adw_2^*) \circ \phi \circ (adw_1) \circ \psi^{-1}$. Then $\hat{\tau} \circ \alpha = \lambda \hat{\tau}$. Therefore $\lambda \in \mathfrak{S}(A)$. $\square$

Example 4.21. Let $\{\lambda_1, \dots, \lambda_n\}$ be nonzero positive numbers such that the closed additive subgroup of $\mathbb{R}$ generated by $\{\lambda_1, \dots, \lambda_n\}$ is $\mathbb{R}$ and $\mathcal{O}_n$ the Cuntz algebra generated by n isometries $S_1, \dots, S_n$. There exists a one-parameter automorphism group $\alpha : \mathbb{R} \rightarrow \text{Aut}(\mathcal{O}_n)$ given by $\alpha_t(S_j) = e^{it\lambda_j} S_j$. Define $A := \mathcal{O}_n \rtimes_{\alpha} \mathbb{R}$. Then A is a simple stable separable C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ and $\mathfrak{S}(A) = \mathbb{R}_+^{\times}$ [20] and [21]. Therefore $\mathcal{F}(A) = \mathbb{R}_+^{\times}$ by the corollary above.

Finally we state a direct relation between the fundamental group of C^* -algebras and that of von Neumann algebras.

Proposition 4.22. Let A be a σ -unital infinite-dimensional simple C^* -algebra with unique (up to scalar multiple) densely defined lower semicontinuous trace τ . Assume that τ is a normalized trace. Consider the GNS representation $\pi_{\tau} : A \rightarrow B(H_{\tau})$ and the associated factor $\pi_{\tau}(A)''$ of type II_1 . Then $\mathcal{F}(A) \subset \mathcal{F}(\pi_{\tau}(A)'')$. In particular, if $\mathcal{F}(\pi_{\tau}(A)'') = \{1\}$, then $\mathcal{F}(A) = \{1\}$.

Proof. Let h be a positive element in $A \otimes \mathbb{K}$ such that A is isomorphic to $\overline{h(A \otimes \mathbb{K})h}$. We denote by $\tilde{\tau}$ the restriction of $\hat{\tau}$ on $\overline{h(A \otimes \mathbb{K})h}$. By the uniqueness of trace, $\pi_{\tau}(A)''$ is isomorphic to $\pi_{\tilde{\tau}}(\overline{h(A \otimes \mathbb{K})h})''$. Define $p := \int_0^{\|h\|} dE_t$ where $\{E_t : 0 \leq t \leq \|h\|\}$ is the spectral projections of $\pi_{\tilde{\tau}}(h)$. Then $d_{\tau}(h) = \hat{\tau}(p)$. A standard argument shows $p\pi_{\tilde{\tau}}(A \otimes \mathbb{K})''p$ is isomorphic to $\pi_{\tilde{\tau}}(\overline{h(A \otimes \mathbb{K})h})''$. Therefore $\mathcal{F}(A) \subset \mathcal{F}(\pi_{\tau}(A)'')$. $\square$

REFERENCES

- [1] B. Blackadar, *Operator Algebras : Theory of C^* -Algebras and von Neumann Algebras*, Encyclopaedia of Mathematical Sciences, **122**, Springer, 2006.
- [2] B. Blackadar and D. Handelman, *Dimension functions and Traces on C^* -algebras*, J. Funct. Anal. **45** (1982), no.3, 297–340.
- [3] B. Blackadar and A. Kumjian, *Skew products of relations and the structure of simple C^* -algebras*, Math. Z. **189** (1985), no. 1, 55–63.
- [4] L. G. Brown, *Stable isomorphism of hereditary subalgebras of C^* -algebras*, Pacific J. Math. **71** (1977), 335–348.
- [5] L. G. Brown, P.Green and M. A. Rieffel, *Stable isomorphism and strong Morita equivalence of C^* -algebras*, Pacific J. Math. **71** (1977), 349–363.
- [6] F. Combes and H. Zettl, *Order structures, traces and weights on Morita equivalent C^* -algebras*, Math. Ann. **265** (1983), no. 1, 67–81.
- [7] A. Connes, *A factor of type II_1 with countable fundamental group*, J. Operator Theory **4** (1980), 151–153.
- [8] J. Cuntz, *Dimension functions on simple C^* -algebras*, Math. Ann. **233** (1978), no. 2, 145–153.
- [9] J. Cuntz and G. K. Pedersen, *Equivalence and traces on C^* -algebras*, J. Funct. Anal. **33** (1979), no. 2, 135–164.
- [10] G. A. Elliott, *An invariant for simple C^* -algebras*, Canadian Mathematical Society. 1945–1995, Vol. 3, 61–90, Canadian Math. Soc., Ottawa, ON, 1996.
- [11] G. A. Elliott and J. Villadsen, *Perforated ordered K_0 -groups*, Canad. J. Math. **52** (2000), no. 6, 1164–1191.
- [12] M. Frank and D. Larson, *A module frame concept for Hilbert C^* -modules*, Contemporary Mathematics **247** (1999), 207–233.

- [13] M. Frank and D. Larson, *Frames in Hilbert C^* -modules and C^* -algebras*, J. Operator Theory **48** (2002), 273–314.
- [14] M. Izumi, T. Kajiwara and Y. Watatani, *KMS states and branched points*, Ergodic Theory Dynam. Systems **27** (2007), no. 6, 1887–1918.
- [15] B. Jacelon, *A simple self-absorbing, stably projectionless C^* -algebra*, preprint, arXiv:1006.5397v1 [math.OA].
- [16] T. Kajiwara, C. Pinzari and Y. Watatani, *Ideal structure and simplicity of the C^* -algebras generated by Hilbert bimodules*, J. Funct. Anal. **159** (1998), no. 2, 295–322.
- [17] T. Kajiwara, C. Pinzari and Y. Watatani, *Jones index theory for Hilbert C^* -bimodules and its equivalence with conjugation theory*, J. Funct. Anal. **215** (2004), no. 1, 1–49.
- [18] T. Kajiwara and Y. Watatani, *Jones index theory by Hilbert C^* -bimodules and K -theory*, Trans. Amer. Math. Soc. **352** (2000), 3429–3472.
- [19] G. G. Kasparov, *Hilbert C^* -modules: theorems of Stinespring and Voiculescu*, J. Operator Theory **4** (1980), no. 1, 133–150.
- [20] A. Kishimoto and A. Kumjian, *Simple stably projectionless C^* -algebras arising as crossed products*, Canad. J. Math. **48** (1996), no. 5, 980–996.
- [21] A. Kishimoto and A. Kumjian, *Crossed products of Cuntz algebras by quasi-free automorphisms*, in Operator algebras and their applications (Waterloo, ON, 1994/1995), 173–192, Fields Inst. Commun., 13, Amer. Math. Soc., Providence, RI, 1997.
- [22] K. Kodaka, *Full projections, equivalence bimodules and automorphisms of stable algebras of unital C^* -algebras*, J. Operator Theory, **37** (1997), 357–369.
- [23] K. Kodaka, *Picard groups of irrational rotation C^* -algebras*, J. London Math. Soc. (2) **56** (1997), 179–188.
- [24] K. Kodaka, *Projections inducing automorphisms of stable UHF-algebras*, Glasg. Math. J. **41** (1999), no. 3, 345–354.
- [25] M. Lacas and S. Neshveyev, *KMS states of quasi-free dynamics on Pimsner algebras*, J. Funct. Anal. **211** (2004), no. 2, 457–482.
- [26] E. C. Lance, *Hilbert C^* -modules*, London Mathematical Society Lecture Note Series, **210**, Cambridge University Press, Cambridge, 1995.
- [27] V. M. Manuilov and E. V. Troitsky, *Hilbert C^* -Modules*, Translations of Mathematical Monographs, **226**, American Mathematical Society, Providence, RI, 2005.
- [28] F. Murray and J. von Neumann, *On rings of operators IV*, Ann. Math. **44**, (1943), 716–808.
- [29] N. Nawata and Y. Watatani, *Fundamental group of simple C^* -algebras with unique trace*, Adv. Math. **225** (2010), no. 1, 307–318.
- [30] N. Nawata and Y. Watatani, *Fundamental group of simple C^* -algebras with unique trace II*, preprint, arXiv:0911.0238v1 [math.OA].
- [31] G. K. Pedersen, *Measure theory for C^* algebras, III*, Math. Scand. **25** (1969), 71–93.
- [32] G. K. Pedersen, *C^* -Algebras and Their Automorphism Groups*, Academic Press, London-New York-San Francisco, 1979.
- [33] C. Pinzari, Y. Watatani and K. Yonetani, *KMS states, entropy and the variational principle in full C^* -dynamical systems*, Comm. Math. Phys. **213** (2000), no. 2, 331–379.
- [34] S. Popa, *Strong rigidity of II_1 factors arising from malleable actions of w -rigid groups, I*, Invent. Math. **165** (2006), 369–408.
- [35] S. Popa and S. Vaes, *Actions of $\mathbb{F}_\infty$ whose II_1 factors and orbit equivalence relations have prescribed fundamental group*, J. Amer. Math. Soc. **23** (2010), 383–403.
- [36] F. Radulescu, *The fundamental group of the von Neumann algebra of a free group with infinitely many generators is $\mathbb{R}_+$* , J. Amer. Math. Soc. **5** (1992), 517–532.
- [37] S. Razak, *On the classification of simple stably projectionless C^* -algebras*, Canad. J. Math. **54** (2002), no. 1, 138–224.
- [38] I. Raeburn and D. P. Williams, *Morita Equivalence and Continuous-Trace C^* -Algebras*, Mathematical Surveys and Monographs, **60**, American Mathematical Society, Providence, RI, 1998.
- [39] M. A. Rieffel, *Morita equivalence for operator algebras*, Operator algebras and applications, Part I (Kingston, Ont., 1980), pp. 285–298, Proc. Sympos. Pure Math., **38**, Amer. Math. Soc., Providence, R.I., 1982.

- [40] K. W. Tsang, *On the positive tracial cones of simple stably projectionless C^* -algebras*, J. Funct. Anal. **227** (2005), no. 1, 188–199.
- [41] D. Voiculescu, *Circular and semicircular systems and free product factors*, in Operator algebras, unitary representations, enveloping algebras, and invariant theory, Progr. Math. 92, Birkhäuser, Boston, 1990, 45–60.
- [42] Y. Watatani, *Index for C^* -subalgebras*, Memoir AMS **424** (1990).

(Norio Nawata) GRADUATE SCHOOL OF MATHEMATICS, KYUSHU UNIVERSITY, MO-
TOOKA, FUKUOKA, 819-0395, JAPAN

E-mail address: `n-nawata@math.kyushu-u.ac.jp`